

## Sheet 1

### Exercise 1.1

#### Part 1

For any  $n \in \mathbb{N}$ , we set  $f_n := \mathbf{1}_{[n, n+1]}$ .

1. Show that for any  $x \in \mathbb{R}_+$ ,  $\lim_{n \rightarrow +\infty} f_n(x) = 0$
2. Show that for any  $n \in \mathbb{N}$ , we have  $\int_{\mathbb{R}_+} f_n(x) dx = 1$

#### Part 2

We will show that the sequence  $(f_n)_{n \in \mathbb{N}}$  does not satisfy the following property: there exist a non-negative function  $g \in L^1(\mathbb{R}_+)$  such that

$$\text{a.e. } x \in \mathbb{R}_+, \forall n \in \mathbb{N}, \quad |f_n(x)| \leq g(x). \quad (1)$$

1. Show that for any  $x \in \mathbb{R}_+$

$$\sup_{n \in \mathbb{N}} \{|f_n(x)|\} = 1.$$

2. Show that, if a measurable function  $g : \mathbb{R}_+ \rightarrow \mathbb{R}$  satisfying (1), then  $g \notin L^1(\mathbb{R}_+)$ .

### Exercise 1.2 (The Fourier transform of complex Gaussians)

Let  $a \in \mathbb{C}$  such that  $\text{Re}(a) > 0$ . The goal of this exercise is to show that

$$\forall x \in \mathbb{R}^d, \quad \int_{\mathbb{R}^d} e^{-ix \cdot \xi} e^{-\frac{|x|^2}{2a}} dx = (2a\pi)^{\frac{d}{2}} e^{-a|\xi|^2} \quad (2)$$

#### Part 1

For any  $x \in \mathbb{R}$ , we define  $h(x) := e^{-\frac{x^2}{2a}}$ . We assume that  $h \in \mathcal{S}(\mathbb{R}^d)$ .

1. Show that  $h'(x) = -\frac{x}{a}h(x)$ .
2. Show that  $h' \in L^1(\mathbb{R}^d)$  and that  $\widehat{h'}(\xi) = i\xi\widehat{h}(\xi)$ .
3. Show that  $\widehat{h'}(\xi) = -i\xi\widehat{h}(\xi)$ .
4. Recall that

$$\int_{\mathbb{R}} h(x) dx = \sqrt{2a\pi}.$$

Show that  $\widehat{h}(0) = \sqrt{2a\pi}$ .

5. Deduce that  $\widehat{h}$  is the solution of the following Cauchy problem

$$\begin{cases} \widehat{h}'(\xi) = -ia\xi\widehat{h}(\xi) & \text{in } \mathbb{R}, \\ \widehat{h}(0) = \sqrt{2a\pi}. \end{cases} \quad (3)$$

6. Deduce from the Cauchy-Lipschitz theorem that, for any  $\xi \in \mathbb{R}$

$$\widehat{h}(\xi) = \sqrt{2\pi a} e^{-a|\xi|}.$$

**Part 2** By remarking that for any  $x = (x_1, \dots, x_d) \in \mathbb{R}^d$  we have

$$e^{-\frac{|x|^2}{2a}} = \prod_{j=1}^d h(x_j),$$

show Formula (2).

### Exercise 1.3 (The heat equation)

Let  $u_0 \in \mathcal{S}(\mathbb{R}^d)$ . For any  $t \geq 0$  and  $\xi \in \mathbb{R}^d$ , we set

$$u(t, x) := \frac{1}{(2\pi)^d} \int_{\mathbb{R}^d} e^{ix \cdot \xi} e^{-t|\xi|^2} \widehat{u}_0(\xi) d\xi.$$

#### Part 1

1. Show that for any  $(t, x) \in (0, +\infty) \times \mathbb{R}^d$ , we have  $\partial_t u(t, x) := \int_{\mathbb{R}^d} (-|\xi|^2) e^{ix \cdot \xi} e^{-t|\xi|^2} \widehat{u}_0(\xi) d\xi$ .
2. Show that  $u \in \mathcal{C}^\infty((0, +\infty) \times \mathbb{R}^d)$ .
3. Show that  $\partial_t u - \Delta u = 0$  in  $(0, +\infty) \times \mathbb{R}^d$ .

#### Part 2

1. Show that

$$\forall (t, x) \in (0, +\infty) \times \mathbb{R}^d, \quad \frac{1}{(2\pi)^d} \int_{\mathbb{R}^d} e^{ix \cdot \xi} e^{-t|\xi|^2} \widehat{u}_0(\xi) d\xi = \frac{1}{(4\pi t)^{\frac{d}{2}}} \int_{\mathbb{R}^d} e^{-\frac{|x-y|^2}{4t}} u_0(y) dy.$$

2. Show that  $\lim_{t \rightarrow 0^+} u(t, x) = u_0(x)$ .
3. Deduce that for any  $x \in \mathbb{R}^d$ , we have  $u(0, x) = u_0(x)$ .

#### Part 3

Show that, for any  $f \in \mathcal{S}(\mathbb{R}^d)$ , we have

$$f(x) = \frac{1}{(2\pi)^d} \int_{\mathbb{R}^d} e^{ix \cdot \xi} \widehat{f}(\xi) d\xi.$$

**Homework (hand in on 22.01.2025).****Exercise 1.4 (The generalised Leibniz rule)**

For multiindices  $\alpha, \beta \in \mathbb{N}^d$ , we declare that  $\beta \leq \alpha$  if  $\beta_j \leq \alpha_j$  for all  $j = 1, \dots, d$ . Denote by

$$\binom{\alpha}{\beta} = \prod_{j=1}^d \binom{\alpha_j}{\beta_j}.$$

Prove the generalised Leibniz formula for  $f, g \in C^{|\alpha|}(\mathbb{R}^d)$

$$\partial^\alpha (fg) = \sum_{\beta \leq \alpha} \binom{\alpha}{\beta} (\partial^\beta f)(\partial^{\alpha-\beta} g).$$

**Exercise 1.5 (The Schrödinger equation)**

Let  $u_0 \in \mathcal{S}(\mathbb{R}^d)$ . For any  $(t, x) \in \mathbb{R} \times \mathbb{R}^d$  we set

$$u(t, x) := \frac{1}{(2\pi)^d} \int_{\mathbb{R}^d} e^{ix \cdot \xi} e^{it|\xi|} \widehat{u}_0(\xi) d\xi.$$

- Show that  $u \in \mathcal{C}^\infty(\mathbb{R} \times \mathbb{R}^d)$ .
- Show that  $u$  solves the Schrödinger equation

$$\begin{cases} \partial_t u + i\Delta u = 0, & \text{in } \mathbb{R} \times \mathbb{R}^d, \\ \lim_{t \rightarrow 0} u(t, x) = u_0(x), & \text{in } \mathbb{R}^d. \end{cases} \quad (4)$$

**Exercise 1.6 (The wave equation)**

Let  $u_0$  and  $u_1$  in  $\mathcal{S}(\mathbb{R}^d)$ . For any  $(t, x) \in \mathbb{R} \times \mathbb{R}^d$  we set

$$u(t, x) := \frac{1}{(2\pi)^d} \int_{\mathbb{R}^d} e^{ix \cdot \xi} \cos(t|\xi|) \widehat{u}_0(\xi) d\xi + \frac{1}{(2\pi)^d} \int_{\mathbb{R}^d} e^{ix \cdot \xi} \frac{\sin(t|\xi|)}{|\xi|} \widehat{u}_1(\xi) d\xi.$$

- Show that  $u \in \mathcal{C}^\infty(\mathbb{R} \times \mathbb{R}^d)$ .
- Show that  $u$  solves the wave equation

$$\begin{cases} \partial_t^2 u - \Delta u = 0, & \text{in } \mathbb{R} \times \mathbb{R}^d, \\ \lim_{t \rightarrow 0} u(t, x) = u_0(x) \text{ and } \lim_{t \rightarrow 0} \partial_t u(t, x) = u_1(x), & \text{in } \mathbb{R}^d. \end{cases} \quad (5)$$